

Three loop $\overline{\text{MS}}$ transversity operator anomalous dimensions for
fixed moment $n \leq 8$

J.A. Gracey,
Theoretical Physics Division,
Department of Mathematical Sciences,
University of Liverpool,
P.O. Box 147,
Liverpool,
L69 3BX,
United Kingdom.

Abstract. We compute the anomalous dimensions of the transversity operator at three loops in the $\overline{\text{MS}}$ scheme for fixed moment n where $n \leq 8$. The results for the RI' renormalization scheme are also provided for an arbitrary linear covariant gauge for $n \leq 7$.

Recently, the three loop anomalous dimension for the unpolarized twist-2 flavour non-singlet and singlet Wilson operators have been determined analytically for all values of the operator moment n , [1, 2, 3, 4]. This was a formidable undertaking spanning ten years and relied on cutting edge computer algebra and symbolic manipulation techniques implemented on high performance computers. With the obvious necessity of such results to ensure the complete two loop evolution of the Wilson coefficients and obtain more precise estimates of quantities such as $\alpha_s(M_Z)$, there is also the need to extend such computations to problems involving spin. Indeed in this area, one quantity which will be of interest is that relating to transversity which was originally introduced in [5, 6, 7]. This corresponds to the probability of finding a quark in a transversely polarized nucleon polarized parallel to the nucleon versus that of the nucleon in the antiparallel polarization. Although experimentally it is harder to extract information on the transversity compared with usual deep inelastic scattering, one will still require the anomalous dimensions of the underlying operator to as high a loop order as is computationally feasible for accurate renormalization group evolution. Currently the two loop anomalous dimensions are available for arbitrary moment, n , in the $\overline{\text{MS}}$ scheme, [8, 9, 10, 11, 12]. However, given the symbolic manipulation machinery now available [1, 2, 3, 4, 13], it is clearly only a matter of time before the full n -dependent three loop $\overline{\text{MS}}$ results are determined. Although the underlying transversity operator differs from that of the Wilson operators of [1, 2, 3, 4], computationally it is on a par with the non-singlet work of [14]. Prior to the full n -dependent results of [1, 2, 3, 4] for the twist-2 Wilson operators, one approach was to carry out a fixed moment determination of the anomalous dimensions. Essentially the first even moments to $n = 16$ (apart from $n = 14$) were determined, [14, 15, 16]. Although eventually superseded by the analytic result, given the huge number of Feynman diagrams to evaluate by tedious recurrence relations, having information on the final results from an *independent* calculation provided an important crosscheck on the full n -dependent expressions. This is particularly the case when computations of gauge invariant quantities are simplified by choosing to work in the Feynman gauge. This substantially reduces the large number of integration by parts and hence the computation time is significantly smaller than, say, in an arbitrary linear covariant gauge. In other words working in a specific gauge means that the internal strong check of observing the gauge parameter cancellation for a gauge independent quantity is crucially absent.

Given the potential determination of analytic anomalous dimensions for the transversity operator in the foreseeable future, it is the purpose of this article to repeat the approach of [14] and provide the $\overline{\text{MS}}$ anomalous dimensions for fixed moments at three loops up to and including moment $n = 8$. This builds on the low moment results of [17, 18] where the anomalous dimensions of the tensor current and second moment were determined. More recently the results for moments $n = 3$ and 4 were provided in [19]. In [18, 19] the *primary* aim was to provide the finite parts of a specific Green's function in order to aid lattice measurements of the same quantity. In particular those measurements had to match onto the ultraviolet part of the Green's function and the provision of the answer at three loops was necessary to help make the extraction of lattice results as precise as possible. However, the work also required performing the renormalization in the lattice renormalization scheme, known as the modified regularization invariant (RI') scheme, [20, 21]. It has a continuum definition which is discussed at length in [22, 17]. As the anomalous dimension of a gauge invariant operator is gauge dependent in a mass dependent scheme, the computation had to be performed in an arbitrary linear covariant gauge. Though for practical reasons the lattice calculations were performed in the Landau gauge. Therefore, as a second thread to this article, we will also determine the transversity anomalous dimensions in the RI' scheme in an arbitrary linear covariant gauge. Though for computational reasons this will be restricted to $n \leq 7$.

First, we discuss the basic properties of the transversity operator and outline our computa-

tional strategy. The operator is defined by, [8, 9, 10],

$$\mathcal{O}^{\mu\nu_1\dots\nu_n} = \mathcal{S}\bar{\psi}\sigma^{\mu\nu_1}D^{\nu_2}\dots D^{\nu_n}\psi \quad (1)$$

where $\sigma^{\mu\nu} = \frac{1}{2}[\gamma^\mu, \gamma^\nu]$ and D_μ is the covariant derivative involving the coupling constant g . The operation $\mathcal{S}$ denotes symmetrization of the Lorentz indices $\{\nu_1, \dots, \nu_n\}$ as well as ensuring that the operator is traceless according to the rules

$$\begin{aligned} \eta_{\mu\nu_i}\mathcal{O}^{\mu\nu_1\dots\nu_i\dots\nu_n} &= 0 \quad (i \geq 2) \\ \eta_{\nu_i\nu_j}\mathcal{O}^{\mu\nu_1\dots\nu_i\dots\nu_j\dots\nu_n} &= 0. \end{aligned} \quad (2)$$

To renormalize (1) we follow a procedure similar to [17, 18] where the operator is inserted at zero momentum into a quark 2-point function, $\langle\psi(p)\mathcal{O}(0)\bar{\psi}(-p)\rangle$, where p is the momentum. In order to apply the MINCER algorithm, [23], written in the symbolic manipulation language FORM, [24, 25], one needs to saturate the Lorentz indices with the appropriate tensor. This is because the MINCER formalism can only be applied to massless three loop Lorentz scalar 2-point Feynman integrals, [23]. Here, since we are not interested in the *finite* part of this Green's function, we merely multiply it by a Lorentz tensor which has the same symmetry and tracelessness properties as the original operator itself. This leads to an immediate algebraic simplification. When the fully symmetrized and traceless operator is inserted into the Green's function, there is a part involving products of the tensor $\eta^{\mu\nu}$. These derive from ensuring the tensor is overall traceless. However, when these terms multiply the projection tensor such terms will give zero. Therefore, in constructing the Feynman rules for the operator insertion for the current calculation, one needs only to consider the part of the operator which is independent of the $\eta^{\mu\nu}$ tensors. In other words, the object $\bar{\psi}\sigma^{\mu(\nu_1}D^{\nu_2}\dots D^{\nu_n)}\psi$. For high moments, this represents a huge reduction in work such as the tedious but automatic derivation of the full Feynman rules, which can involve a significantly large number of terms. Instead the main work is in the construction of the projection tensor. However, this is achieved computer algebraically by writing down the complete set of independent objects built from one $\sigma^{\mu\nu}$ tensor, together with the appropriate numbers of $\eta^{\mu\nu}$ tensors and momenta p^σ such that the number of free Lorentz indices equates with that of the original operator. These independent tensors are then symmetrized automatically with respect to the indices $\{\nu_1, \dots, \nu_n\}$ and the arbitrary coefficients chosen so that the overall projection tensor is traceless according to (2). As was indicated in [18, 19], there are three independent projections and therefore for the determination of the anomalous dimensions one needs only to select one of these for the projection procedure. It is preferable to choose the most algebraically compact one to minimize computation time. We note that this procedure is in contrast to the fixed moment strategy of [14] where the null vector Δ_μ was introduced. However, as was discussed in [14, 26], one needs to handle the resulting integrals using a different projection technique. Finally, we note that our procedure automatically ensures that there is no operator mixing, [19]. The Feynman diagrams required for $\langle\psi(p)\mathcal{O}(0)\bar{\psi}(-p)\rangle$ are generated automatically with the QGRAF package, [27], and converted to FORM input notation prior to the application of the three loop MINCER algorithm. There are 3 one loop, 37 two loop and 684 three loop Feynman diagrams and throughout we have used dimensional regularization in $d = 4 - 2\epsilon$ dimensions.

We now record that the three loop $\overline{\text{MS}}$ transversity anomalous dimensions, $\gamma^{(n)}(a)$, for $n = 5, 6$ and 7 are

$$\begin{aligned} \gamma^{(5)}(a) &= \frac{92}{15}C_F a + [189515C_A - 41674C_F - 79810T_F N_f] \frac{C_F a^2}{6750} \\ &+ \left[(190836000\zeta(3) + 1975309075) C_A^2 \right. \\ &\quad \left. - (572508000\zeta(3) + 325464235) C_A C_F \right] \end{aligned}$$

$$\begin{aligned}
& - (1192320000\zeta(3) + 511395100) C_A T_F N_f \\
& + (381672000\zeta(3) - 254723696) C_F^2 \\
& + (1192320000\zeta(3) - 989903260) C_F T_F N_f \\
& - 83718800 T_F^2 N_f^2 \Big] \frac{C_F a^3}{12150000} + O(a^4)
\end{aligned} \tag{3}$$

$$\begin{aligned}
\gamma^{(6)}(a) = & \frac{34}{5} C_F a + [204770 C_A - 42129 C_F - 88810 T_F N_f] \frac{C_F a^2}{6750} \\
& + \Big[(707616000\zeta(3) + 7527909825) C_A^2 \\
& - (2122848000\zeta(3) + 1373507730) C_A C_F \\
& - (4626720000\zeta(3) + 1841332000) C_A T_F N_f \\
& + (1415232000\zeta(3) - 684744816) C_F^2 \\
& + (4626720000\zeta(3) - 3910683210) C_F T_F N_f \\
& - 320975800 T_F^2 N_f^2 \Big] \frac{C_F a^3}{42525000} + O(a^4)
\end{aligned} \tag{4}$$

and

$$\begin{aligned}
\gamma^{(7)}(a) = & \frac{258}{35} C_F a + [75266555 C_A - 15484767 C_F - 33149830 T_F N_f] \frac{C_F a^2}{2315250} \\
& + \Big[(3517994592000\zeta(3) + 38365845513450) C_A^2 \\
& - (10553983776000\zeta(3) + 5978407701105) C_A C_F \\
& - (24084527040000\zeta(3) + 9039144860900) C_A T_F N_f \\
& + (7035989184000\zeta(3) - 4192441946262) C_F^2 \\
& + (24084527040000\zeta(3) - 20698675427220) C_F T_F N_f \\
& - 1651311191600 T_F^2 N_f^2 \Big] \frac{C_F a^3}{204205050000} + O(a^4)
\end{aligned} \tag{5}$$

where $a = g^2/(16\pi^2)$, $\zeta(n)$ is the Riemann zeta function, N_f is the number of quark flavours and the colour group Casimirs are given by

$$\text{Tr} \left(T^a T^b \right) = T_F \delta^{ab} \ , \ T^a T^a = C_F I \ , \ f^{acd} f^{bcd} = C_A \delta^{ab} \tag{6}$$

and T^a are the generators for the colour group with structure constants f^{abc} . We have also calculated the $n = 8$ moment but modified the method used to find (3), (4) and (5). In order to reduce the computation time we performed the calculation in the Feynman gauge. Whilst this significantly reduces the number of integration by parts needed to be carried out by the MINCER algorithm, for example, the check of gauge invariance of the final result in the $\overline{\text{MS}}$ scheme is removed. However, as a partial check on the automatic construction of the $n = 8$ Feynman rules and projection tensor, we have performed the two loop calculation in an arbitrary linear covariant gauge and reconstructed the gauge independent result of [8, 9, 10, 11, 12]. Thus, at three loops in the $\overline{\text{MS}}$ scheme we have

$$\begin{aligned}
\gamma^{(8)}(a) = & \frac{551}{70} C_F a + [1270588235 C_A - 251839827 C_F - 568470280 T_F N_f] \frac{C_F a^2}{37044000} \\
& + \Big[(57548150352000\zeta(3) + 651153115163775) C_A^2 \\
& - (172644451056000\zeta(3) + 110529966326535) C_A C_F
\end{aligned}$$

$$\begin{aligned}
& - (411490679040000\zeta(3) + 148380288276500) C_A T_F N_f \\
& + (115096300704000\zeta(3) - 55777651074312) C_F^2 \\
& + (411490679040000\zeta(3) - 356756758487220) C_F T_F N_f \\
& - 27902636165600 T_F^2 N_f^2 \Big] \frac{C_F a^3}{3267280800000} + O(a^4) .
\end{aligned} \tag{7}$$

As further checks on our results, (3), (4), (5) and (7), we note that for $\overline{\text{MS}}$ all the two loop expressions agree with those of [8, 9, 10, 11, 12], when they are evaluated for specific n . Moreover, (3), (4) and (5) are clearly independent of the linear covariant gauge fixing parameter, α . Further, the coefficients of the leading order large N_f term at three loops, corresponding to the $C_F T_F^2 N_f^2$ term, agrees with the analytic evaluation at $O(1/N_f)$ for arbitrary n given in [18]. A final check is provided by the fact that the triple and double poles in ϵ of the resulting operator renormalization constant are reproduced precisely in agreement with the renormalization group equation prediction for all four cases. This is important since we followed the procedure of [28] for renormalizing operators in automatic symbolic manipulation programmes. For the specific case of the Lie group $SU(3)$ we have

$$\begin{aligned}
\gamma^{(5)}(a) \Big|^{SU(3)} &= \frac{368}{45} a - 2 [119715 N_f - 1538939] \frac{a^2}{30375} \\
&- \left[188367300 N_f^2 + 8942400000 \zeta(3) N_f + 12843253410 N_f \right. \\
&\quad \left. - 954180000 \zeta(3) - 144207743479 \right] \frac{a^3}{82012500} + O(a^4)
\end{aligned} \tag{8}$$

$$\begin{aligned}
\gamma^{(6)}(a) \Big|^{SU(3)} &= \frac{136}{15} a - 2 [44405 N_f - 558138] \frac{a^2}{10125} \\
&- \left[240731850 N_f^2 + 11566800000 \zeta(3) N_f + 16107360420 N_f \right. \\
&\quad \left. - 1179360000 \zeta(3) - 183119500163 \right] \frac{a^3}{95681250} + O(a^4)
\end{aligned} \tag{9}$$

$$\begin{aligned}
\gamma^{(7)}(a) \Big|^{SU(3)} &= \frac{344}{35} a - 2 [16574915 N_f - 205153309] \frac{a^2}{3472875} \\
&- \left[206413898950 N_f^2 + 10035219600000 \zeta(3) N_f \right. \\
&\quad + 13678917121415 N_f - 977220720000 \zeta(3) \\
&\quad \left. - 156962874344971 \right] \frac{a^3}{76576893750} + O(a^4)
\end{aligned} \tag{10}$$

and

$$\begin{aligned}
\gamma^{(8)}(a) \Big|^{SU(3)} &= \frac{1102}{105} a - [284235140 N_f - 3475978269] \frac{a^2}{27783000} \\
&- \left[20926977124200 N_f^2 + 1028726697600000 \zeta(3) N_f \right. \\
&\quad + 1381224814218690 N_f - 95913583920000 \zeta(3) \\
&\quad \left. - 15957293707773841 \right] \frac{a^3}{7351381800000} + O(a^4) .
\end{aligned} \tag{11}$$

where $C_F = 4/3$, $C_A = 3$ and $T_F = 1/2$.

Finally, we note that the results analogous to (3), (4) and (5) in the RI' scheme are

$$\begin{aligned}
\gamma_{\text{RI}'}^{(5)}(a) = & \frac{92}{15} C_F a \\
& + \left[(30825\alpha^2 + 92475\alpha + 1740690) C_A - 166696 C_F - 676560 T_F N_f \right] \frac{C_F a^2}{27000} \\
& + \left[(194197500\alpha^4 + 1854279000\alpha^3 - 583200000\zeta(3)\alpha^2 + 8993896875\alpha^2 \right. \\
& \quad - 6026400000\zeta(3)\alpha + 30074295375\alpha - 37353312000\zeta(3) \\
& \quad + 356401468700) C_A^2 \\
& + (91239750\alpha^3 - 209956950\alpha^2 - 4997987400\alpha + 1076976000\zeta(3) \\
& \quad - 60979980560) C_A C_F \\
& - (1726200000\alpha^2 - 1555200000\zeta(3)\alpha + 10041363000\alpha + 17858880000\zeta(3) \\
& \quad + 253330505600) C_A T_F N_f \\
& + (1289803200\alpha + 27164160000\zeta(3) - 22363266560) C_F T_F N_f \\
& \left. + (10686816000\zeta(3) - 7132263488) C_F^2 + 41629683200 T_F^2 N_f^2 \right] \frac{C_F a^3}{340200000} \\
& + O(a^4)
\end{aligned} \tag{12}$$

$$\begin{aligned}
\gamma_{\text{RI}'}^{(6)}(a) = & \frac{34}{5} C_F a \\
& + \left[(33075\alpha^2 + 99225\alpha + 1957910) C_A - 168516 C_F - 769360 T_F N_f \right] \frac{C_F a^2}{27000} \\
& + \left[(119070000\alpha^4 + 1160028000\alpha^3 - 364500000\zeta(3)\alpha^2 + 5621954625\alpha^2 \right. \\
& \quad - 3766500000\zeta(3)\alpha + 18833983125\alpha - 22639824000\zeta(3) \\
& \quad + 231206558900) C_A^2 \\
& + (49437000\alpha^3 - 176477400\alpha^2 - 3356146800\alpha - 1150848000\zeta(3) \\
& \quad - 40247461840) C_A C_F \\
& - (1058400000\alpha^2 - 972000000\zeta(3)\alpha + 6258495000\alpha + 11741760000\zeta(3) \\
& \quad + 165502535600) C_A T_F N_f \\
& + (866102400\alpha + 18040320000\zeta(3) - 14470193920) C_F T_F N_f \\
& \left. + (6469632000\zeta(3) - 3130262016) C_F^2 + 27562342400 T_F^2 N_f^2 \right] \frac{C_F a^3}{194400000} \\
& + O(a^4)
\end{aligned} \tag{13}$$

and

$$\begin{aligned}
\gamma_{\text{RI}'}^{(7)}(a) = & \frac{258}{35} C_F a \\
& + \left[(12006225\alpha^2 + 36018675\alpha + 739917710) C_A \right. \\
& \quad \left. - 61939068 C_F - 292181680 T_F N_f \right] \frac{C_F a^2}{9261000} \\
& + \left[(1588423567500\alpha^4 + 15732651627000\alpha^3 - 4900921200000\zeta(3)\alpha^2 \right. \\
& \quad + 76165633191375\alpha^2 - 50642852400000\zeta(3)\alpha + 255327460003875\alpha \\
& \quad \left. - 301268627856000\zeta(3) + 3230428196219700) C_A^2 \right.
\end{aligned}$$

$$\begin{aligned}
& + \left(598161044250\alpha^3 - 2827913095350\alpha^2 - 47764761023700\alpha \right. \\
& \quad \left. - 30807568512000\zeta(3) - 558295101833000 \right) C_A C_F \\
& - \left(14119320600000\alpha^2 - 13069123200000\zeta(3)\alpha + 84542124471000\alpha \right. \\
& \quad \left. + 164110847040000\zeta(3) + 2326270368265200 \right) C_A T_F N_f \\
& + (12326389941600\alpha + 254163329280000\zeta(3) - 198131779069280) C_F T_F N_f \\
& + (84431870208000\zeta(3) - 50309303355144) C_F^2 \\
& + 391057633881600 T_F^2 N_f^2 \Big] \frac{C_F a^3}{2450460600000} + O(a^4) . \tag{14}
\end{aligned}$$

To summarize, this renormalization scheme is defined by ensuring that the *finite* part of the renormalized Green's function $\langle \psi(p) \mathcal{O}(0) \bar{\psi}(-p) \rangle$, multiplied by the projector, is given purely by its tree value only, [20, 21, 17, 18, 19]. Further, it is important to note that in (12), (13) and (14) the variables a and α are to be regarded as RI' quantities. The relation to the $\overline{\text{MS}}$ variables are given to three loops in [17] for an arbitrary linear covariant gauge. Since our main motivation is to provide the $\overline{\text{MS}}$ anomalous dimensions, the $n = 8$ moment in the RI' scheme is clearly not available since the corresponding $\overline{\text{MS}}$ computation was restricted to the Feynman gauge.

We conclude by noting that we have provided higher moments of the transversity operator at three loops in both the $\overline{\text{MS}}$ and RI' schemes. Together with the earlier results of [20, 21, 18, 19], there are now eight fixed moment $\overline{\text{MS}}$ anomalous dimensions available prior to an explicit n -dependent computation. This is similar to the situation with regard to the flavour non-singlet twist-2 operator where there were seven fixed moment anomalous dimensions available prior to the provision of the full n -dependent expression. Whilst it is still in principle possible to compute even higher moments using the method we have discussed here, we believe we have reached a computational limit beyond which it is not viable to proceed. For instance, for the three loop $n = 8$ Feynman gauge calculation, the necessary Feynman rules took around 36 hours to be generated electronically on a dual opteron 64 bit SMP machine (2 GHz), resulting in just under 5.5×10^6 terms. The former is an order of magnitude larger than the time required for the $n = 7$ case.

Acknowledgements. The author thanks Dr P.E.L. Rakow and Dr C. McNeile for valuable discussions.

References.

- [1] S. Moch, J.A.M. Vermaseren & A. Vogt, Nucl. Phys. **B688** (2004), 101.
- [2] S. Moch, J.A.M. Vermaseren & A. Vogt, Nucl. Phys. **B691** (2004), 129.
- [3] S. Moch, J.A.M. Vermaseren & A. Vogt, Nucl. Phys. Proc. Suppl. **135** (2004), 137.
- [4] S. Moch, J.A.M. Vermaseren & A. Vogt, Phys. Lett. **B606** (2005), 123.
- [5] J.P. Ralston & D.E. Soper, Nucl. Phys. **B152** (1979), 109.
- [6] R.L. Jaffe & X. Ji, Phys. Rev. Lett. **67** (1991), 552; Nucl. Phys. **B375** (1992), 527.
- [7] J.I. Cortes, B. Pire & J.P. Ralston, Z. Phys. **C55** (1992), 409.
- [8] F. Baldracchini, N.S. Craigie, V. Roberto & M. Socolovsky, Fortschr. Phys. **30** (1981), 505.

- [9] X. Artru & M. Mekhfi, Z. Phys. **C45** (1990), 669.
- [10] A. Hayashigaki, Y. Kanazawa & Y. Koike, Phys. Rev. **D56** (1997), 7350.
- [11] W. Vogelsang, Phys. Rev. **D57** (1998), 1886.
- [12] J. Blümlein, Eur. Phys. J. **C20** (2001), 683.
- [13] J.A.M. Vermaseren, Int. J. Mod. Phys. **A14** (1999), 2037.
- [14] S.A. Larin, T. van Ritbergen & J.A.M. Vermaseren, Nucl. Phys. **B427** (1994), 41.
- [15] A. Rétey & J.A.M. Vermaseren, Nucl. Phys. **B604** (2001), 281.
- [16] J. Blümlein & J.A.M. Vermaseren, Phys. Lett. **B606** (2005), 130.
- [17] J.A. Gracey, Nucl. Phys. **B662** (2003), 247.
- [18] J.A. Gracey, Nucl. Phys. **B667** (2003), 242.
- [19] J.A. Gracey, JHEP **0610** (2006), 040.
- [20] G. Martinelli, C. Pittori, C.T. Sachrajda, M. Testa & A. Vladikas, Nucl. Phys. **B445** (1995), 81.
- [21] E. Franco & V. Lubicz, Nucl. Phys. **B531** (1998), 641
- [22] K.G. Chetyrkin & A. Rétey, Nucl. Phys. **B583** (2000), 3.
- [23] S.G. Gorishny, S.A. Larin, L.R. Surguladze & F.K. Tkachov, Comput. Phys. Commun. **55** (1989), 381.
- [24] S.A. Larin, F.V. Tkachov & J.A.M. Vermaseren, “The Form version of Mincer”, NIKHEF-H-91-18.
- [25] J.A.M. Vermaseren, math-ph/0010025.
- [26] S.A. Larin, P. Nogueira, T. van Ritbergen & J.A.M. Vermaseren, Nucl. Phys. **B492** (1997), 338.
- [27] P. Nogueira, J. Comput. Phys. **105** (1993), 279.
- [28] S.A. Larin & J.A.M. Vermaseren, Phys. Lett. **B303** (1993), 334.